

Competitive Monte Carlo methods for the Pricing of Asian Options

B. Lapeyre* E. Temam †

August 24, 2000

Abstract

We explain how a carefully chosen scheme can lead to competitive Monte Carlo algorithm for the computation of the price of Asian options. We give evidence of the efficiency of these algorithms with a mathematical study of the rate of convergence and a numerical comparison with some existing methods.

KEY WORDS: Asian option, Monte Carlo methods, Numerical methods, Diffusion process.

1 Introduction

Monte Carlo methods are known to be useful when the state dimension is large. This is widely true but we will give here an example of a small dimension problem coming from finance where a Monte Carlo (helped by a variance reduction technique) can be more efficient than other known methods.

This example is based on the price of an Asian option (see subsection 2.1). This problem is known to be computationally hard and a lot of literature deals with this problem: using either analytic methods ([10], [9]), numerical methods based on the partial differential equation associated ([4], [7], [12], [16]) or Monte Carlo methods ([13]). The originality of this work is to propose (and to give precise results) new time schemes approximation for the integral of the Black & Scholes model: this point is really a source of concern when using Monte Carlo methods as noted by Madan, Fu and Wang ([9], page 14). We will show that, when we use a suitable scheme and variance reduction, a Monte Carlo method can be more competitive than other methods under some circumstances.

In order to get precise and complete numerical results, we have undertaken extensive comparisons with most of the other known methods : Forward Shooting Grid ([2]), Hull and White ([11]), Finite Difference and already quoted Monte Carlo methods. For moderate precision 10^{-2} , tree methods (FSG and Hull and White) are the most

*CERMICS, Ecole Nationale des Ponts et Chaussées, 6 et 8 avenue Blaise Pascal, 77455 Marne La Vallée, - FRANCE, e-mail : bl@cermics.enpc.fr

†Université Paris VI, Laboratoire de Probabilités et Modèles Aléatoires and CERMICS, Ecole Nationale des Ponts et Chaussées, 6 et 8 avenue Blaise Pascal, 77455 Marne La Vallée, - FRANCE, e-mail : temam@cermics.enpc.fr

efficient. But, surprisingly, really large precision (10^{-4}) can only be reached by Monte Carlo method with relatively large time step (~ 1 month).

This article is organized in the following way: we first introduce the mathematical context, then we present the Monte Carlo schemes in sections 1-2, section 3 contains the proof of their convergence in the L^p spaces and section 4 quickly describes the variance reduction method used in this article. The end of the paper is devoted to numerical tests and comparisons.

2 Mathematical Context

2.1 Financial Background

To describe the price of an asset at time t , we use the Black and Scholes model with a risky asset (a share of price S_t at time t) and a no-risk asset (whose price is S_t^0 at time t). The price S_t is given by the following stochastic differential equation

$$dS_t = S_t(\mu dt + \sigma dB_t),$$

where μ and σ are two positive constants and (B_t) is a standard Brownian motion. The price of the no-risk asset satisfies the ordinary differential equation

$$dS_t^0 = rS_t^0 dt.$$

As usual, we introduce the process $W_t = B_t + \frac{\mu-r}{\sigma}t$ which is a Brownian motion under an adapted probability, called the neutral risk probability and denoted by \mathbb{P} . Thus, the risky asset satisfies a new stochastic differential equation

$$dS_t = S_t(rt + \sigma dW_t),$$

whose solution is

$$S_t = S_{T_0} \exp\left(\sigma W_t - \frac{\sigma^2}{2}t + rt\right).$$

S_{T_0} is the price of the asset at the beginning of the modeling.

Asian options (or options on average) is the general name for a class of options whose payoff depends of the mean of the price of the risky asset on a given period. Thus, the price of an Asian option with maturity T can be written:

$$V(t, S, A) = e^{-r(T-t)} \mathbb{E}f(S_t, A_S(T_0, t)).$$

where

$$A_S(T_0, t) = \frac{1}{t - T_0} \int_{T_0}^t S_u du.$$

From now on, we choose $T_0 = 0$, with no lost of generality. The function f depends of the type of the options

- For a call with fixed strike: $f(s, a) = (a - K)_+$

- For a put with fixed strike: $f(s, a) = (K - a)_+$
- For a call with floating strike: $f(s, a) = (s - a)_+$
- For a put with floating strike: $f(s, a) = (a - s)_+$

2.2 Practical Schemes

If we want to use Monte Carlo methods to compute a price, we have to simulate the average of S_t , therefore we need to approximate an integral. Here, it is not necessary to approximate S_t because it can be exactly simulated. With this aim in view, the interval $[0, T]$ will be divided into N steps. The step size will be noted $h = T/N$, and we define the times $t_k = kT/N = kh$.

The standard scheme We introduce three schemes to estimate $Y_T = \int_0^T S_u du$. Since we are able to simulate S_t at a given t , the integral can be approached by Riemann sums:

$$\mathbf{Y}_T^{r,n} = \mathbf{h} \sum_{k=0}^{n-1} \mathbf{S}_{t_k}. \quad (2.1)$$

For example, if M denote the number of drawing in the Monte Carlo method, an approximation of the price at maturity of a fixed strike Asian call is given by

$$\frac{e^{-rT}}{M} \sum_{j=1}^M \left(\frac{h}{T} \sum_{k=0}^{n-1} S_{t_k} - K \right)_+.$$

Note that the time complexity of this algorithm is $O\left(\frac{1}{NM}\right)$ (this is true for every kind of Monte Carlo methods) and that it involves two kinds of errors: the Monte Carlo error and the time step error. The Monte Carlo error is of order $\frac{\sigma}{\sqrt{M}}$ and the time step error is harder to evaluate (see proposition 3.3).

The scheme (2.1) can be interpreted as the second variable in the Euler approximation for the following stochastic differential equation

$$dU_t = B(U_t)dt + \Sigma(U_t)dW_t \quad \text{with } U_t = \begin{bmatrix} S_t \\ Y_t \end{bmatrix}, \quad B(U_t) = \begin{bmatrix} rS_t \\ S_t \end{bmatrix} \quad \text{and } \Sigma(U_t) = \begin{bmatrix} \sigma(S_t) \\ 0 \end{bmatrix}$$

Higher accuracy schemes One way to obtain higher accuracy for the integral approximation is to remark that in L^2 , the closest random variable to $\left(\frac{1}{T} \int_0^T S_s ds - K\right)_+$ when the $(S_{t_k}, k = 0, \dots, N)$ are known is given by:

$$\mathbb{E} \left(\left(\frac{1}{T} \int_0^T S_u du - K \right)_+ \middle| \mathcal{B}_h \right), \quad (2.2)$$

with \mathcal{B}_h is the σ -field generated by the $(S_{t_k}, k = 0, \dots, N)$. Of course, it is theoretically impossible to compute explicitly this conditional expectation (it is certainly harder than

to obtain an explicit formula for V). But, since the conditional law of W_u with respect to \mathcal{B}_h for $u \in [t_k, t_{k+1}]$ is given by

$$\mathcal{L}(W_u \mid W_{t_k} = x, W_{t_{k+1}} = y) = \mathcal{N}\left(\frac{t_{k+1} - u}{h}x + \frac{u - t_k}{h}y, \frac{(t_{k+1} - u)(u - t_k)}{h}\right), \quad (2.3)$$

one can in principle compute

$$\left(\mathbb{E}\left(\frac{1}{T}\int_0^T S_u du \mid \mathcal{B}_h\right) - K\right)_+ = \left(\frac{1}{T}\int_0^T \mathbb{E}\left(S_u \mid \mathcal{B}_h\right) du - K\right)_+ \quad (2.4)$$

as a function of $(W_{t_k}, k = 0, \dots, N)$. Jensen inequality proves that (2.4) is smaller than (2.2), but we will see that (2.2) is already a really good approximation of Y_T (see proposition 3.3). Using the law described by (2.3), we get

$$\begin{aligned} & \mathbb{E}\left[\frac{1}{T}\int_0^T S_u du \mid \mathcal{B}_h\right] \\ &= \frac{1}{T}\sum_{k=0}^{n-1}\int_{t_k}^{t_{k+1}} e^{(r-\frac{\sigma^2}{2})u} e^{\sigma\frac{t_{k+1}-u}{h}W_{t_k} + \sigma\frac{u-t_k}{h}W_{t_{k+1}} + \frac{\sigma^2}{2}\frac{(t_{k+1}-u)(u-t_k)}{h}} du \\ &= \frac{1}{T}\sum_{k=0}^{n-1}\int_{t_k}^{t_{k+1}} e^{\sigma\frac{u-t_k}{h}(W_{t_{k+1}}-W_{t_k}) - \frac{\sigma^2}{2}\frac{(u-t_k)^2}{h} + ru} e^{\sigma W_{t_k} - \frac{\sigma^2}{2}t_k} du \end{aligned}$$

In Monte Carlo simulation, this approximation is used in a double loop (in times and in the number of simulation), so it is really necessary to simplify this formula. Hence, a formal Taylor expansion can be done with h small, which leads to a more practical scheme:

$$\mathbf{Y}_T^{e,n} = \frac{h}{T}\sum_{k=0}^{n-1} \mathbf{S}_{t_k} \left(1 + \frac{rh}{2} + \sigma\frac{\mathbf{W}_{t_{k+1}} - \mathbf{W}_{t_k}}{2}\right). \quad (2.5)$$

Remark 2.1. Note that this scheme is equivalent to the well known trapezoidal method. We will prove that,

$$\mathbb{E}\left(Y_T^{e,n} - \frac{1}{T}\sum_{k=0}^{n-1} h\frac{S_{t_k} + S_{t_{k+1}}}{2}\right)^2 = O\left(\frac{1}{n^3}\right).$$

So, since the rate of convergence of (2.5) is in $1/n$ (see proposition 3.3), the error above is negligible.

Proof. This result can be obtained using a Taylor expansion:

$$\begin{aligned} \frac{1}{T}\sum_{k=0}^{n-1} h\frac{S_{t_k} + S_{t_{k+1}}}{2} &= \frac{1}{T}\sum_{k=0}^{n-1} \frac{hS_{t_k}}{2} (e^{\sigma(W_{t_{k+1}}-W_{t_k}) - \frac{\sigma^2}{2}h + rh} + 1) \\ &= \frac{1}{T}\sum_{k=0}^{n-1} \frac{hS_{t_k}}{2} (2 + \sigma(W_{t_{k+1}} - W_{t_k}) + rh) + \dots \end{aligned}$$

which is exactly the scheme (2.5). The term of order $\sigma^2 h$ and the quadratic variation of $\sigma(W_{t_{k+1}} - W_{t_k})$ cancel each other out, so we does not retain it. The mathematical proof of this remark is the same as proposition 3.3. \square

Our last scheme is quite similar. As the Brownian motion is a Gaussian process, $\int_0^T W_u du$ has a normal density w.r.t. the Lebesgues measure on \mathbb{R} and can easily be simulated. Hence,

$$\begin{aligned} Y_T &= \frac{1}{T} \int_0^T S_u du \\ &= \frac{1}{T} \sum_{k=0}^{n-1} S_{t_k} \int_{t_k}^{t_{k+1}} e^{\sigma(W_u - W_{t_k}) - \frac{\sigma^2}{2}(u - t_k) + r(u - t_k)} du. \end{aligned}$$

So, using a Taylor expansion again, we obtain the scheme:

$$\mathbf{Y}_T^{\mathbf{p},n} = \frac{1}{T} \sum_{k=0}^{n-1} \mathbf{S}_{t_k} \left(\mathbf{h} + \frac{\mathbf{r}h^2}{2} + \sigma \int_{t_k}^{t_{k+1}} (\mathbf{W}_u - \mathbf{W}_{t_k}) d\mathbf{u} \right). \quad (2.6)$$

Remark 2.2. In practice, to simulate this scheme, we have, at each time step, to simulate $W_{t_{k+1}}$ knowing W_{t_k} and $(\int_{t_k}^{t_{k+1}} W_u du \mid W_{t_k}, W_{t_{k+1}})$. For the second random variable, we use the law (2.3) and for the first one the fact that $(W_{t_{k+1}} - W_{t_k}, k = 0, \dots, N - 1)$ are Gaussian i.i.d. variables.

Remark 2.3. This scheme can be generalized easily to a higher class of diffusion process. Let S_t be a diffusion with a drift $b(S_t)$ and a diffusion term $\sigma(S_t)$, and $Y_t = \int S_t dt$. One can use the Euler scheme S_t^n to simulate S_t and

$$Y_T^n = \sum_{k=0}^{n-1} S_{t_k}^n \left(h + \int_{t_k}^{t_{k+1}} (S_s^n - S_{t_k}^n) ds \right).$$

Note that when the diffusion S_t can be directly simulated, the following scheme is sufficient

$$Y_T^n = \sum_{k=0}^{n-1} S_{t_k} \left(h + \int_{t_k}^{t_{k+1}} (S_s^n - S_{t_k}) ds \right),$$

We can show that the weak convergence holds at the rate $1/n^{3/2}$.

We will now prove some results on the speed of convergence of these 3 schemes. The weak convergence can be obtained for the scheme (2.1) (at the rate $1/n$) and (2.6) (at the rate $1/n^{3/2}$) under a weak assumption on f using Malliavin calculus and techniques developed by Bally and Talay ([1]). We refer to [17] for a complete proof of these convergence results.

3 Convergence in the L^p spaces

In this section, we are interested by the strong convergence of the schemes. In L^p , one can only have some upper bound of the rate of convergence, but the precise expansion can only be obtained in the particular case $p = 2$.

Let us start with two well known but important results.

Proposition 3.1. *For a Black and Scholes diffusion,*

$$\mathbb{E}|S_t - S_s|^{2q} \leq C_q |t - s|^q$$

This proposition is available for any diffusion with Lipschitz coefficient (see [15]).

The following will also be useful (see [6] chapter 3 for a proof):

Lemma 3.1. *Let $Z_t = Z_0 + \int_0^t A_s dW_s + \int_0^t B_s ds$ where B_s is a vector in \mathbb{R}^n , A_s a matrix in $\mathbb{R}^{n \times d}$, and W_t a d dimensional Brownian motion. (Z_t is an Itô process so A and B are adapted, $\int |A_s| ds < +\infty$ and $\mathbb{E} \int B_s^2 ds < +\infty$).*

Then, Z_t satisfies

$$\mathbb{E}|Z_t|^p \leq \mathbb{E}|Z_0|^p + C \int_0^t \mathbb{E}(|Z_s|^p + |A_s|^p + |B_s|^p) ds$$

We can now obtain precise results for the rate of convergence of our schemes.

Proposition 3.2. *With the above notations, there exist three non decreasing maps $K_1(T)$, $K_2(T)$, $K_3(T)$ such that,*

$$\left(\mathbb{E} \left(\sup_{t \in [0, T]} |Y_t^{r, n} - Y_t|^{2q} \right) \right)^{\frac{1}{2q}} \leq \frac{K_1(T)}{n} \quad (3.1)$$

$$\left(\mathbb{E} \left(\sup_{t \in [0, T]} |Y_t^{e, n} - Y_t|^{2q} \right) \right)^{\frac{1}{2q}} \leq \frac{K_2(T)}{n} \quad (3.2)$$

$$\left(\mathbb{E} \left(\sup_{t \in [0, T]} |Y_t^{p, n} - Y_t|^{2q} \right) \right)^{\frac{1}{2q}} \leq \frac{K_3(T)}{n^{3/2}} \quad (3.3)$$

Proof. Let us begin with the inequality (3.1). If we note $\epsilon_t = Y_t^{r, n} - Y_t$, one has

$$\begin{aligned} \forall t \in [t_k, t_{k+1}] \quad \epsilon_t &= \epsilon_{t_k} + \int_{t_k}^t (S_s - S_{t_k}) ds \\ &= \epsilon_{t_k} + \int_{t_k}^t (t - u) r S_u du + \int_{t_k}^t (t - u) \sigma S_u dW_u \end{aligned}$$

So, the lemma 3.1 implies that

$$\forall t \in [t_k, t_{k+1}] \quad \mathbb{E}|\epsilon_t|^{2q} \leq (\mathbb{E}|\epsilon_{t_k}|^{2q} + h^{2q+1}) + \int_{t_k}^t \mathbb{E}|\epsilon_s|^{2q} ds.$$

Applying Gronwall's lemma, we get that

$$\mathbb{E}|\epsilon_{t_{k+1}}|^{2q} \leq (\mathbb{E}|\epsilon_{t_k}|^{2q} + h^{2q+1})e^h$$

Since $x_{n+1} \leq ax_n + b$ implies $x_n \leq a^n x_0 + ne^{n(a-1)}b$ for $a \geq 1$, we have

$$\mathbb{E}|\epsilon_{t_k}|^{2q} \leq Ch^{2q}$$

We conclude with the inequality of Burkholder-Davis-Gundy. We use the same kind of arguments for the two other schemes writing them as:

$$\begin{aligned} Y_t - Y_t^{e,n} &= Y_{t_k} - Y_{t_k}^{e,n} + \int_{t_k}^t \int_{t_k}^s r(S_u - S_{t_k}) du ds + \int_{t_k}^t \int_{t_k}^s \sigma(S_u - \frac{h}{2}S_{t_k}) dW_u ds \\ &= Y_{t_k} - Y_{t_k}^{e,n} + \int_{t_k}^t r(t-u)(S_u - S_{t_k}) du + \int_{t_k}^t \sigma((t-u)S_u - \frac{h}{2}S_{t_k}) dW_u \end{aligned}$$

and,

$$\begin{aligned} Y_t - Y_t^{p,n} &= Y_{t_k} - Y_{t_k}^{p,n} + \int_{t_k}^t \int_{t_k}^s r(S_u - S_{t_k}) du ds + \int_{t_k}^t \int_{t_k}^s \sigma(S_u - S_{t_k}) dW_u ds \\ &= Y_{t_k} - Y_{t_k}^{e,n} + \int_{t_k}^t r(t-u)(S_u - S_{t_k}) du + \int_{t_k}^t \sigma(t-u)(S_u - S_{t_k}) dW_u. \end{aligned}$$

From this point, it is easy to get the inequalities (3.2) and (3.3). □

These inequalities give an upper bound for the rate of convergence, but we can obtain more precise results. Moreover, if we study the L^2 convergence, we can obtain an expansion, and this proves the exact rate of convergence.

Proposition 3.3. *With the above notations and assumptions, it holds that*

$$\sqrt{\mathbb{E} \left[\frac{1}{T} \int_0^T S_u du - \mathbb{E} \left(\frac{1}{T} \int_0^T S_u du \mid \mathcal{B}_h \right) \right]^2} = \frac{\sigma}{n} \sqrt{\frac{e^{(\sigma^2+2r)T} - 1}{12(\sigma^2 + 2r)}} + O\left(\frac{1}{n\sqrt{n}}\right) \quad (3.4)$$

$$\sqrt{\mathbb{E} \left[\frac{1}{T} \int_0^T S_u du - Y_T^{e,n} \right]^2} = \frac{\sigma}{n} \sqrt{\frac{e^{(\sigma^2+2r)T} - 1}{12(\sigma^2 + 2r)}} + O\left(\frac{1}{n\sqrt{n}}\right) \quad (3.5)$$

$$\sqrt{\mathbb{E} \left[\frac{1}{T} \int_0^T S_u du - Y_T^{r,n} \right]^2} = \frac{K_2}{n} + O\left(\frac{1}{n\sqrt{n}}\right) \quad \text{where } K_2 \geq \sigma \sqrt{\frac{e^{(\sigma^2+2r)T} - 1}{12(\sigma^2 + 2r)}} \quad (3.6)$$

$$\sqrt{\mathbb{E} \left[\frac{1}{T} \int_0^T S_u du - Y_T^{p,n} \right]^2} = \frac{1}{n\sqrt{n}} \sqrt{T\sigma^4 \frac{e^{(\sigma^2+2r)T} - 1}{12(\sigma^2 + 2r)}} + O\left(\frac{1}{n^2}\right). \quad (3.7)$$

Remark that the same rate of convergence occurs in equations (3.4) and (3.5), and furthermore the first term in the expansion is the same. It implies that we did not lose anything by approximating the conditional expectation with the scheme (2.5).

If we are interested in the strong convergence, it seems that the most efficient scheme is the last one. We also remark that the schemes (2.1) and (2.5) have the same rate of convergence and that the first terms in the expansions are quite similar.

The proof of these 4 equalities are similar and we will treat only the last one.

Proof of (3.7) Let us note $A_{u,t_k} = e^{\sigma(W_u - W_{t_k}) - \frac{\sigma^2}{2}(u-t_k) + r(u-t_k)} - 1 - r(u-t_k) - \sigma(W_u - W_{t_k})$. Then we get,

$$\begin{aligned} \mathbb{E} \left[\frac{1}{T} \int_0^T S_u du - Y_T^{p,n} \right]^2 &= \frac{2}{T^2} \sum_{0 \leq j < i \leq N-1} \int_{t_i}^{t_{i+1}} dv \int_{t_j}^{t_{j+1}} du \mathbb{E}(A_{v,t_i} A_{u,t_j} S_{t_i} S_{t_j}) \\ &\quad + \frac{2}{T^2} \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} dv \int_{t_k}^u du \mathbb{E}(A_{u,t_k} A_{v,t_k} S_{t_k}^2). \end{aligned}$$

Since A_{v,t_i} is independent of \mathcal{F}_{t_i} , it follows that

$$\begin{aligned} \mathbb{E}(A_{v,t_i} A_{u,t_j} S_{t_i} S_{t_j}) &= e^{(\sigma^2+r)t_j + rt_i} (e^{r(v-t_i)} - 1 - r(v-t_i)) \\ &\quad (e^{(r+\sigma^2)(u-t_i)} - 1 - (r+\sigma^2)(u-t_i)). \end{aligned}$$

With a normalization of the variable in the integrand we get,

$$\begin{aligned} \frac{2}{T^2} \sum_{0 \leq j < i \leq N-1} \int_{t_i}^{t_{i+1}} dv \int_{t_j}^{t_{j+1}} du \mathbb{E}(A_{v,t_i} A_{u,t_j} S_{t_i} S_{t_j}) &= \frac{2}{T^2} \left(\sum_{0 \leq j < i \leq N-1} e^{(\sigma^2+r)t_j + rt_i} \right) h^2 \int_0^1 dv \int_0^1 du \\ &\quad (e^{rhv} - 1 - rhv)(e^{(r+\sigma^2)(hu)} - 1 - (r+\sigma^2)hu). \end{aligned}$$

It is clear that $\left(\sum_{0 \leq j < i \leq N-1} e^{(\sigma^2+r)t_j + rt_i} \right) = O\left(\frac{1}{h^2}\right)$, which leads to

$$\frac{2}{T^2} \sum_{0 \leq j < i \leq N-1} \int_{t_i}^{t_{i+1}} dv \int_{t_j}^{t_{j+1}} du \mathbb{E}(A_{v,t_i} A_{u,t_j} S_{t_i} S_{t_j}) = O(h^4).$$

We use the same method for the second term,

$$\begin{aligned} \mathbb{E}(A_{u,t_k} A_{v,t_k} S_{t_k}^2) &= e^{(\sigma^2+2r)t_k} \left(- (1 + r(u-t_k))(e^{r(v-t_k)} - 1 - r(v-t_k)) \right. \\ &\quad \left. + e^{r(u-t_k)}(e^{(\sigma^2+r)(v-t_k)} - 1 - (r+\sigma^2)(v-t_k)) + \sigma^2(v-t_k)(e^{r(v-t_k)} - 1) \right). \end{aligned}$$

Consequently,

$$\begin{aligned} \frac{2}{T^2} \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} dv \int_{t_k}^u du \mathbb{E}(A_{u,t_k} A_{v,t_k} S_{t_k}^2) &= \frac{2}{T^2} \left(\sum_{k=0}^{N-1} e^{(\sigma^2+2r)t_k} \right) h^2 \int_0^1 du \int_0^u dv \left(\frac{\sigma^4 h^2 v^2}{2} + O(h^3) \right) \\ &= h^3 \frac{\sigma^4 e^{(\sigma^2+2r)T} - 1}{12(\sigma^2 + 2r)}. \end{aligned}$$

4 Variance Reduction

To increase the efficiency of the Monte Carlo simulation a variance reduction method can be used. We follow the method developed by Kemna and Vorst [13]. It consists in approximating $\frac{1}{T} \int_0^T S_u du$ by $\exp\left(\frac{1}{T} \int_0^T \log(S_u) du\right)$. We can expect that these two random variables are similar since r and σ are not too large.

The random variable $Z' = \frac{1}{T} \int_0^T \log(S_u) du$ obviously has a normal law, and so we can compute explicitly

$$\mathbb{E}(e^{-rT} (\exp(Z') - K)_+).$$

As a consequence, we can choose the random variable Z defined by

$$Z = e^{-rT} (xe^{(r-\frac{\sigma^2}{2})\frac{T}{2} + \frac{\sigma}{T} \int_0^T W_u du} - K)_+,$$

as our control variable.

Note that the control variable has to be computed with the path of the Brownian motion already simulated. Consequently, each control variable has to be adapted to the schemes. So, we retain the same approximation for $\int_0^T W_u du$ as for $\int_0^T S_u du$:

$$\begin{aligned} \text{For (2.1)} \quad Z_T^{r,n} &= e^{-rT} (xe^{(r-\frac{\sigma^2}{2})\frac{T}{2} + \frac{\sigma}{T} \sum_{k=0}^{n-1} hW_{t_k}} - K)_+ \\ \text{For (2.5)} \quad Z_T^{e,n} &= e^{-rT} (xe^{(r-\frac{\sigma^2}{2})\frac{T}{2} + \frac{\sigma}{T} \sum_{k=0}^{n-1} \frac{h}{2}(W_{t_k} + W_{t_{k+1}})} - K)_+ \\ \text{For (2.6)} \quad Z_T^{p,n} &= e^{-rT} (xe^{(r-\frac{\sigma^2}{2})\frac{T}{2} + \frac{\sigma}{T} \sum_{k=0}^{n-1} \int_{t_k}^{t_{k+1}} W_u du} - K)_+ \end{aligned}$$

Note that, as it is explained in remark 2.2, when we use the scheme (2.6), we simulate at each step $W_{t_{k+1}}$ then the random variables ξ_k defined by $(\int_{t_k}^{t_{k+1}} W_u du \mid W_{t_k}, W_{t_{k+1}})$. By

writing $\sum_{k=0}^{n-1} \int_{t_k}^{t_{k+1}} W_u du$ in the last equation above, we intend the sum of the generated variables i.e. $\sum_{k=0}^{n-1} \xi_k$.

5 Numerical computation

In this section we compare the accuracy of various schemes to compute the price of an Asian option. First, we will study the convergence of our schemes with and without the variance reduction method. This leads to the foreseeable conclusion that the variance reduction is essential to the efficiency of our schemes. In a second part, we compare the Monte Carlo methods with other methods of pricing, namely Forward Shooting Grid, Hull & White and a finite difference method.

In the graphics or table of this section, the letter R represents the scheme (2.1), E the scheme (2.5) and BB the scheme (2.6).

5.1 Convergence of the Monte Carlo Methods

5.1.1 Simulation without variance reduction technique

We test the three schemes of subsection 2.2 with the following values: $T = 1$ year, $r = 0.1$, $X_0/K = 1$ and $\sigma = 0.2$.

The first simulation is made without any variance reduction technique and with 100000 Monte Carlo loops which is already high. The results presented in the table 1, show that even if the convergence seems to be fast, the confidence intervals are too large for a real financial use (penny accuracy is required).

Scheme	Number of time step	Price	Confidence Intervals at 95%
(2.1)	10	6.396	[6.348,6.444]
	50	6.978	[6.926,7.030]
	100	6.930	[6.877,6.982]
(2.5)	10	6.997	[6.944,7.050]
	50	7.034	[6.981,7.087]
	100	7.037	[6.984,7.090]
(2.6)	10	7.065	[7.012,7.118]
	50	7.059	[7.006,7.112]
	100	7.055	[7.002,7.108]

Table 1: Confidence Intervals versus time step number when no variance reduction technique is used.

As we will see later, the real price of this call is not far from 7.04. The table 1 shows that the first scheme is not very efficient compared to the others. The surprise is that (2.5) and (2.6) seems to have the same rate of convergence whereas the proposition 3.3 ensures that the second one is more efficient. But, the justification of the numerical simulation can be based on weak convergence and if we study that kind of convergence, the two schemes will have the same expansion (see [17] for details).

The table 1 shows the limit of the Monte Carlo method. In fact, since our objective is to study the convergence of our schemes when the time step number goes to infinity, we want to avoid problems related to the Monte Carlo approximation. With this aim in view, we have to increase the number of paths which also increases the time of computation dramatically. Variance reduction technique is necessary.

5.1.2 Simulation with variance reduction method

Graphic 1 represents the three schemes computed with the variance reduction technique of section 4 and with the same parameters as above except for the number of loops for the Monte Carlo method which is now 20000 (80 % less than before). We observe again that the schemes (2.5) and (2.6) are quite similar and that the Riemann approximation is less efficient.

The figure 2 which is zoom of the figure 1 with only the two new schemes seems to show that these schemes have the same rate of convergence.




Figure 1: Convergence of the schemes with variance reduction technique; N is the number of time step; $T = 1 \text{ year}$, $r = 0.1$, $X_0/K = 1$ and $\sigma = 0.2$




Figure 2: Convergence of the schemes with variance reduction technique; N is the number of time step; $T = 1 \text{ year}$, $r = 0.1$, $X_0/K = 1$ and $\sigma = 0.2$

Scheme	Number of time step	Price	Confidence Intervals at 95%
(2.1)	10	6.778	[6.776,6.781]
	50	6.991	[6.988,6.995]
	100	7.016	[7.013,7.020]
(2.5)	10	7.035	[7.031,7.038]
	50	7.039	[7.035,7.042]
	100	7.042	[7.038,7.045]
(2.6)	10	7.041	[7.038,7.044]
	50	7.041	[7.037,7.044]
	100	7.041	[7.037,7.044]

Table 2: Confidence Intervals crossed with time step number with variance reduction technique

Scheme	Precision (in %)	Number of time step	Time of computation
(2.1)	5	2	<1s
	1	5	<1s
	0.5	9	1s
	0.1	42	2s
	0.07	65	3s
	0.05	92	3s
	0.03	175	7s
	0.01	800	28s
(2.5)	5	2	< 1s
	1	2	< 1s
	0.5	2	< 1s
	0.1	2	< 1s
	0.07	3	<1s
	0.05	3	<1s
	0.03	4	< 1s
	0.01	9	1s
(2.6)	5	2	<1s
	1	2	<1s
	0.5	2	<1s
	0.1	2	<1s
	0.07	2	<1s
	0.05	2	<1s
	0.03	3	<1s
	0.01	8	1s

Table 3: Minimal number of time steps and time of computation to obtain a given precision

Table 2 represents the dependences between the price of the option and the number of time steps.

Results of table 2 show that the variance reduction method makes the scheme more efficient, both for the convergence and the confidence intervals:

Note that

- The scheme using conditional expectation is a negatively biased approximation.
- The graph show that the time step number which is necessary to have a good precision is less with the schemes (2.5) and (2.6) than with (2.1). As a consequence, the use of low discrepancy sequences can be considered to increase efficiency of our schemes.

The table 3 gives the number of time steps which is necessary to have the confidence interval at 95% include in intervals given by the real value of the option and the precision on the second column. To be able to get a real price, we use zero strike Asian option, and the same other parameters as above (in particular 20000 sample paths).

The results show again how it is efficient to use one of the two new approximations for the integral. Even to get a high precision 0.01%, the schemes (2.5) and (2.6) do not take more than 1 second against 28 seconds for the Riemann sums.

5.2 A numerical study of the rate of convergence




Figure 3: Convergence of the schemes with variance reduction technique; we cross the logarithm of the precision (11.524956 is the real price) with the logarithm of the time step number N

We are trying, in this section, to get numerically the rate of convergence in law of the schemes described in section 2.2. To achieve this, we compute the schemes with a very high number of paths in the Monte Carlo method (10^7) in order to avoid Monte Carlo noise.

The parameters of the option computed are: $r = 0.09$, $\sigma = 0.2$, $T = .4$ and $K = S_{T_0} = 100$.

By running the routine FSG (see [2]) which will be described in the next section with a low time step we have a very good approximation of the real price (~ 11.524956). In figure 3, we plot the logarithm of the difference between this price and the result of the schemes (2.5) and (2.6) as a function of $\log(N)$.

The coefficient of correlation for these two sets of data are -1.46 for (2.6) and -1.33 for (2.5), which approximatively confirms that the rate of weak convergence of (2.5) and (2.6) is $\frac{1}{N^{3/2}}$.

5.3 Comparison with other numerical methods

In this part, we have implemented some other methods commonly used for pricing Asian options:

- A method based on the one dimensional partial derivative equation of which V is solution (see Deynne & Wilmott [4], Ingersoll [12] or Rogers & Shi [16]). This equation is

$$\frac{\partial V}{\partial t}(x, t) + \frac{1}{2}\sigma^2 x^2 \frac{\partial^2 V}{\partial x^2}(x, t) - (\rho(t) + rx) \frac{\partial V}{\partial x}(x, t) = 0 \quad (5.1)$$

with limit condition and Price

$$V(x, T) = x_- \quad Price = S_{T_0} V\left(\frac{K}{S_{T_0}}, 0\right)$$

for a fixed strike and

$$V(x, T) = (1 + x)_- \quad Price = S_{T_0} V(0, 0)$$

for a floating strike. This equation was approximated by a finite difference method with viscosity arguments to increase the efficiency of the method.

- Two methods (often called “tree” methods) based on approximation of the transition kernel of S with the Cox, Ross, Rubinstein discrete model. These methods have been developed by Barraquand and Pudet (FSG of Forward Shooting Grid, [2]) and Hull & White ([11]). We choose for the time step number $N = 12$ (one month), 24, 52 (one week), 100.
- The three schemes for Monte Carlo methods described here, with and without variance reduction techniques. The number of paths was 20000 and the time step numbers $N = 2, 6, 12, 24, 52$

To compare the various schemes, we use the methodology proposed by Broadie et Detemple ([3]). It consists in simulating 200 options, whose parameters are randomly chosen as follows

- The volatility σ follows a uniform law between 0.1 et 0.6.




Figure 4: Results of the advanced test; FSG and Hull stands for the tree methods and MC denotes Monte Carlo methods; KV describes whether or not the variance reduction method is used

- The maturity T is, with probability 0.75, uniform in $\{k/12, k = 1, \dots, 12\}$ and, with probability 0.25 uniform in $\{0, 1, 2, 3, 4, 5\}$.
- The interest rate r is, with probability 0.8, uniform in $[0.0, 0.1]$, and, with probability 0.2 equal to 0.
- S_0 , the initial value of the model (the stock), is uniform between 70 and 130.

The computation has been made on PC with two processor Intel Pentium II at 450Mhz. To be able to evaluate the precision, we chose $K = 0$, which is the only case where there exists a closed formula. The results of this test are given in graphic 4, which represent the logarithm of the mean error

$$E = \sqrt{\frac{1}{200} \sum_{i=1}^{200} \left(\frac{True\ Price - Simulated\ Price}{True\ Price} \right)^2},$$

versus the logarithm of the CPU time of computation.

Graphic 4 can be read as follows

- MC represents Monte Carlo Methods.
- As above, R is the abbreviation for the scheme (2.1), E for (2.5) and BB for (2.6).
- KV is added when the variance reduction method describes above was used.
- FSG is "Forward Shooting Grid Method".
- FD represents the Finite Difference method implemented here.

The results concerning the analytic methods (developed by Geman & Yor [10] and Madan, al. [9]) are unconfirmed. These methods are difficult to implement because of the repetition of numerical inversions of Laplace transform. For the moment their results are disappointing. Using standard algorithms to solve this problem leads to disappointing results. Nevertheless, we are not absolutely sure to have used the best available method.

A standard finite difference on the one dimensional equation (5.1) failed if the volatility is low. Since the variance is a decreasing map of the volatility σ , Monte Carlo methods can be very useful in this case.

We still have to consider the Monte Carlo methods and the Tree methods. For the first class, variance reduction technique and precise schemes ((2.5) et (2.6)) represent the only way to obtain correct results.

To obtain moderate accuracy ($\sim 10^{-3}$), the algorithm of Hull and White seems to be the most efficient. With a low step number (say 6 per year), it gives approximations with a precision at 10^{-3} in a time lower than 0.03 seconds whereas for the same precision, the forward shooting grid takes 0.1 seconds and Monte Carlo methods 0.3 seconds (ten times more!).

At the opposite, if higher precision is required, the Monte Carlo methods become more efficient. In fact, the time complexity of the trees methods is higher (we can see that on the graphic, by comparing the slope of the trees methods and Monte Carlo methods).

For a precision at 10^{-4} , the scheme MC_E_KV runs in 0.4 seconds whereas the algorithm FSG takes 1.3 seconds and Hull et White does not reach this precision for a time step number lower than 100.

The ends of the lines representing the schemes MC_BB_KV and MC_E_KV show that with a time step number above 24, the error due to Monte Carlo becomes greater than the error due to the time step in average.

In conclusion, if we use both the new schemes and the variance reduction technique, Monte Carlo methods can be competitive with the trees methods for pricing Asian options and represent the only way to get high accuracy.

References

- [1] V. Bally and D. Talay. The law of the euler scheme for stochastic differential equations. i: Convergence rate of the distribution function. *Probab. Theory Relat. Fields* 104, No.1, 43-60, 1996.
- [2] J. Barraquand and T. Pudet. Pricing of american path-dependant contingent claims. *Mathematical Finance*, 6:17–51, 1996.
- [3] M. Broadie and J. Detemple. Recent advances in numerical methods for pricing derivative securities. In L. C. G. (ed.) et al. Rogers, editor, *Numerical methods in finance*, pages 43–66, 1997.
- [4] J. Dewynne and P. Wilmott. Asian options as linear complementary problems: Analysis and finite difference solutions. *Advances in Futures and Operations Research* 8, 145-173, 1995.
- [5] D. Dufresne. Laguerre series for asian and other options. *forthcoming in Mathematical Finance*, 1999.
- [6] O. Faure. *Simulation du mouvement brownien et des diffusions*. PhD thesis, Ecole Nationale des Ponts et Chaussées, 1992.
- [7] P.A. Forsyth, K. Vetzal, and R. Zvan. Robust numerical methods for pde models of asian options. Technical report, University of Waterloo, Canada, 1996.
- [8] P.A. Forsyth, K. Vetzal, and R. Zvan. Convergence of lattice and pde methods for pricing asian options. Technical report, University of Waterloo, Canada, 1999.
- [9] M. Fu, D. Madan, and T. Wang. Pricing continous time asian options: A comparison of analytical and monte carlo methods. *forthcoming in the Journal of Computational Finance*, 1996.
- [10] H. Geman and M. Yor. Bessel processes, asian options, and perpetuities. *Mathematical Finance*, 3(4):349–375, 1993.
- [11] J. Hull and A. White. Efficient procedures for valuing european and american path dependant options. *Journal of Derivatives*, 1:21–31, 1993.

- [12] J. Ingersoll, Jr. *Theory of Financial Decision Making*. Roman & Littlefield, Totowa, New Jersey, 1987.
- [13] A.G.Z. Kemna and A.C.F. Vorst. A pricing method for options based on average asset values. *J. Banking Finan.*, March 1990.
- [14] T. Klassen. Simple, fast, and flexible pricing of asian options. *forthcoming in the Journal of Computational Finance*, 2000.
- [15] D. Revuz and M. Yor. *Continuous Martingales and Brownian Motion*. Springer-Verlag, 1994.
- [16] L.C.G. Rogers and Z. Shi. The value of an asian option. *J. Appl. Probab.* 32, No.4, 1077-1088, 1995.
- [17] E. Temam. Monte carlo methods for asian options. Technical Report 144, CER-MICS - Ecole Nationale des Ponts et Chaussées, 1998.